



## Radiochemical analysis for decommissioning of nuclear facilities

Hou, Xiaolin

*Published in:*

9th International Conference on Nuclear and Radiochemistry - NRC9

*Publication date:*

2016

*Document Version*

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*

Hou, X. (2016). Radiochemical analysis for decommissioning of nuclear facilities. In *9th International Conference on Nuclear and Radiochemistry - NRC9: Abstracts* (pp. 89-90)

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## Radiochemical analysis for decommissioning of nuclear facilities

Xiaolin Hou

*Technical University of Denmark, Center for Nuclear Technologies (DTU Nutech), Risø Campus,  
DK-4000 Roskilde, Denmark  
xiho@dtu.dk*

With increasing numbers of nuclear facilities, especially nuclear power reactors, being closed in recent years and from now on, a considerable work is going to be carried out all over the world for decommissioning these nuclear facilities. For this purpose, characterization of various types of materials from the nuclear facilities is required for evaluation of the radioactivity inventory and distribution in various materials and decision making for management of the produced waste. This is implemented by quantitative determination of various radionuclides presenting in these materials.

The neutron activation products of components and impurity in the materials used in the nuclear facilities, such as  $^3\text{H}$ ,  $^{14}\text{C}$ ,  $^{36}\text{Cl}$ ,  $^{41}\text{Ca}$ ,  $^{60}\text{Co}$ ,  $^{55}\text{Fe}$ ,  $^{63}\text{Ni}$ ,  $^{79}\text{Se}$ ,  $^{93}\text{Mo}$ ,  $^{93}\text{Zr}$ ,  $^{94}\text{Nb}$ ,  $^{126}\text{Sn}$ ,  $^{133}\text{Ba}$ ,  $^{152}\text{Eu}$ ,  $^{154}\text{Eu}$  and some transuranics (isotopes of Pu, Np, Am and Cm) are the main contributors to the total radioactivity, especially in the construction materials. While the long-lived fission products, such as  $^{90}\text{Sr}$ ,  $^{99}\text{Tc}$ ,  $^{129}\text{I}$ ,  $^{135}\text{Cs}$  and  $^{137}\text{Cs}$ , are the major concern for materials contaminated by leaked spent nuclear fuel. Of these radionuclides, the gamma emitting radionuclides, such as  $^{60}\text{Co}$ ,  $^{133}\text{Ba}$ ,  $^{152}\text{Eu}$ ,  $^{154}\text{Eu}$  and  $^{137}\text{Cs}$ , are easily measured by gamma spectrometry. While the determination of pure beta and alpha emitters including  $^3\text{H}$ ,  $^{14}\text{C}$ ,  $^{36}\text{Cl}$ ,  $^{41}\text{Ca}$ ,  $^{55}\text{Fe}$ ,  $^{63}\text{Ni}$ ,  $^{90}\text{Sr}$ ,  $^{99}\text{Tc}$ ,  $^{129}\text{I}$  and some transuranics is the major challenges, because they could not be measured without separation from the matrix of the samples and from all other radionuclides, this entitles them as the radionuclides of hard to measure.

Radiochemical analysis is the only way to implement determination of these hard to measure radionuclides by including a separation of individual radionuclides from matrix and other radionuclides before measurement. Although plenty of analytical methods have been reported for the determination of these radionuclides since the discovery of radioactivity, the suitable methods are not always available for the purpose of characterization of various materials from decommissioning. This is attributed to a few reasons including complicate and various components of the sample matrix, some orders of magnitude lower concentration of some target radionuclides comparing to the major radionuclides which needs a high decontamination factors, complicated and active chemical properties of some important target radionuclides, and a large number of samples required to be analyzed during the decommissioning, which needs simple and rapid methods to provide a good analytical capacity. In particular, the sample matrix varies very much from concrete, graphite, exchange resin, to various metals, which requires different radiochemical methods for different sample matrix and target radionuclides.

Besides the conventional radiometric methods for the measurement of radionuclides, such as liquid scintillation counting, low level beta counting, and alpha spectrometry for these hard to measure radionuclides, some new methods have been proposed and applied. Among them, mass spectrometry especially ICP-MS and accelerator mass spectrometry (AMS) has been widely used for the measurement of long-lived radionuclides, such as  $^{41}\text{Ca}$ ,  $^{36}\text{Cl}$ ,  $^{99}\text{Tc}$ ,  $^{129}\text{I}$ ,  $^{135}\text{Cs}$  and transuranics. With these techniques, the measurement time can be significantly reduced from few days to some minutes, enabling to rapid measurement of large number of samples.

In the past few years, a large numbers of radiochemical analytical methods aiming to characterize various decommissioning waste by determination of various radionuclides of hard to measure have been developed in many laboratories. This work aims to present the state of the art analytical methods for characterization of nuclear waste from the decommissioning of nuclear facilities, not only the updated chemical separation

procedure, but also advanced measurement techniques. Meanwhile, some examples of analytical methods will be also presented, including (1) rapid determination of  $^3\text{H}$  and  $^{14}\text{C}$  in solid materials; (2) determination of  $^{36}\text{Cl}$  and  $^{129}\text{I}$  in graphite, concrete and metals (3) Simultaneously determination of multi-nuclides such as  $^{41}\text{Ca}$ ,  $^{55}\text{Fe}$ ,  $^{63}\text{Ni}$ ,  $^{90}\text{Sr}$  and transuranics; (4) Challenge on the determination of low level  $^{93}\text{Mo}$ ,  $^{93}\text{Zr}$ ,  $^{135}\text{Cs}$ ,  $^{79}\text{Se}$  and  $^{126}\text{Sn}$  in decommissioning waste.

### References:

- Hou X.L., Frøsig Østergaard L.; Nielsen S.P., Determination of Ni-63 and Fe-55 in nuclear waste and environmental samples, *Anal. Chim. Acta*, 2005, 535:297-307
- Hou X.L. Determination of C-14 and H-3 in Reactor Graphite and Concrete for Decommission, *Applied Radiation and Isotopes*, 2005, 62:871-882.
- Hou X.L., Radiochemical determination of  $^{41}\text{Ca}$  in reactor concrete for decommissioning, *Radiochim., Acta*, 2005, 93:611-617.
- Hou X.L., Østergaard L.F., Nielsen S.P., Determination of  $^{36}\text{Cl}$  in Nuclear Waste from Reactor Decommissioning, *Anal. Chem.*, 2007, 79, 3126-3134.
- Hou X.L, Radiochemical analysis of radionuclides difficult to measure, *J. Radioanal. Nucl. Chem.*, 2007, 273, 43-48.
- Hou X.L., Roos P., Critical Comparison of Radiometric and Mass Spectrometric Methods for the Determination of Radionuclides in Environmental, Biological and Nuclear Waste samples, *Anal. Chim. Acta*, 2008, 608, 105-139.
- Hou X.L. Determination of long-lived radionuclides in the environment using ICP-MS and AMS, *J. Anal. Sci. Technol.* 2011, 2(Suppl A) A120-124.
- Qiao J.X., Shi K.L., Hou X.L., Nielsen S.P., Roos P., Rapid multi-sample analysis for simultaneous determination of anthropogenic radionuclides in marine environment, *Environment. Sci. Technol.* 2014, 48:3935-3942.
- Shi K., Hou X.L., Roos P., Wu W.S. Stability of technetium and decontamination of Ru and Mo in determination of  $^{99}\text{Tc}$  in environmental solid samples by ICP-MS, *Anal. Chem.*, 2012, 84:2009-2016.
- Zheng J., Bu W.T., Tagami K., Shikamori Y., Nakano K., Uchida S., Ishii N. Determination of  $^{135}\text{Cs}$  and  $^{135}\text{Cs}/^{137}\text{Cs}$  atomic ratio in environmental samples by combining ammonium molybdophosphate (AMP)-selective Cs adsorption and ion-exchange chromatographic separation to triple-quadrupole inductively coupled plasma-mass spectrometry, *Anal. Chem.*, 2014, 86:7103-7110.